

# FINITE SIZE EFFECTS FOR THE DILUTE COUPLING DERRIDA MODEL

large Allakhverdyan A.E., Saakian D.B.

*Yerevan Physics Institute*

Alikhanian Brothers St.2, Yerevan 375036, Armenia

Saakian @ vx1.YERPHI.AM

## Abstract

We consider paramagnetic, spin-glass and ferromagnetic phases. At  $T = 0$  model gives for the some values of connectivity (near the critical) extremal suppression of finite size effects (decoding error probability).

**1. Introduction** Derridas's model [1] is the simplest among the spin-glasses. It has solved in the first level of replica symmetry breaking [2].

May be this is the origin of unique feature-condition for the appearance of ferromagnetic phase coincides with the constraint of Shannon for the optimal coding.

This hypothesis [3] have proved for the general case in [4],[5]

The physical meaning of codes in the approach of Sourlas is this:

The  $N$  numbers  $\xi_i$  are given, taking values  $\pm 1$ .

By means of them we construct  $Z$  numbers  $\tau_i (Z \geq N)$ , then formulate some function of  $N$  discrete variables  $\sigma_i$

$$H(\sigma_i, \tau_j), \quad (1)$$

which has as constants  $Z$  couplings  $\tau_j$ ,

There is a restriction, that single global minimum is configuration

$$\sigma_i = \xi_i \quad (2)$$

It is a simple to construct such function, but then in our problem appears noise. Our couplings  $\tau_i$  with probability  $\frac{1+m}{2}$  conserve their sign, and with probability  $\frac{1-m}{2}$  change

it. Again there is a restriction, that configuration (2) still is a single vacuum for the noisy couplings  $\tau_i$ .

It is possible to do this procedure only for the some (good!) choice of function  $H$ . For this case information theory gives restriction [6]

$$Z[\ln 2 - h(m)] \geq N \ln 2 \quad (3)$$

where for the entropy we have

$$h(m) = -\frac{1+m}{2} \ln \frac{1+m}{2} - \frac{1-m}{2} \ln \frac{1-m}{2} \quad (4)$$

Really our condition (single vacuum) is correct not with probability 1, but a little less:

$$1 - a \exp[-E(m, N/Z)Z] \quad (5)$$

The second member is just the decoding error probability.

On the language of statistical physics we will investigate magnification at  $T = 0$

$$< \xi_i \sigma_j > \quad (6)$$

This expression gives probability, that our condition is full. So decoding error probability is equivalent to finite size effect in the expression of magnetization

In such manner we have coded our original information- $N$  numbers  $\xi_i$ , to noisy number  $\tau_j$ . We can rederive original values of  $N$  numbers  $\xi_i$ , searching (by Monte-Karlo) minimum of function  $H(\sigma_i, \tau_j)$ ,

We want extremely suppress decoding error probability.

The information theory gives restriction for the maximal value of function  $E(m, R)$ , where

$$R = N/Z \quad (7)$$

is a rate of information transmission.

On the [7] finite size effects have calculation at  $T = 0$  for the full connectively case in Derrida model, which corresponds to limit

$$Z/N \rightarrow \infty \quad m \rightarrow 1/2 \quad (8)$$

On the [8] has investigated fine structure of ferromagnetic phase, and has found some subphases, which differs each-others by finite size effects.

In this work we are going to calculate finite size effects for paramagnetic, spin-glass and ferromagnetic phase. For that purposes we are using methods of [9].

## 1 Derivation of REM with weak connectivity

Let us consider hamiltonian

$$H = - \sum_{(i_1 \dots i_p)} \tau_{i_1 \dots i_p} \sigma_{i_1} \dots \sigma_{i_p} \quad (9)$$

where spins  $\tau_i$ , are quenched couplings  $\tau_i$ , taking values  $\pm 1$ ,

In our hamiltonian there are  $C_N^P$  different choice of  $(i_1 \dots i_p)$  and consequently  $C_N^P$  different couplings  $\tau_{i_1 \dots i_p}$ . We consider restriction

$$\sum_{(i_1 \dots i_p)} (\tau_{i_1 \dots i_p})^2 = \alpha N \quad (10)$$

In the bounds of this restriction, which means an weak correlation, all  $\tau_j$  are distributed independently, with the same probabilities

$$\tau_{i_1 \dots i_p} = 0 \quad \text{with probability} \quad 1 - \alpha N / C_N^P \quad (11)$$

$$\tau_{i_1 \dots i_p} = \pm 1 \quad \text{with probability} \quad \frac{1 \pm m}{2} \frac{\alpha N}{C_N^P} \quad (12)$$

On [9] authors considered only conditions (11),(12). Our restriction (10) simplifies calculations. As a results finite size effects are changed only in paramagnetic phase

Let us consider, as in [9], common distribution of  $M$  energy levels.

We have  $M$  configurations with the values of spins  $\sigma_i^\alpha, \leq \alpha \leq M$  and values of energy  $E_\alpha$ .

Let as define

$$P(E_1 \dots E_M) = \langle \delta(E_1 - H(\sigma_i^1)) \dots \delta(E_M - H(\sigma_i^M)) \rangle \quad (13)$$

If it would be possible to calculate. Case with

$$M = 2^N \quad (14)$$

Then we could calculate

$$\langle \ln Z(\tau, T) \rangle = \int \prod_{\alpha=1}^{2^N} dE_{\alpha} P(E_1 \cdots E_{2^N}) \ln \sum_{\alpha=1}^{2^N} \exp(-BE_{\alpha}) \quad (15)$$

We have derived (15), using the fact that our system has only  $2^N$  levels of energy.

That's why we have considered  $Z$  as a function of energy levels  $E_{\alpha}$  (instead of-as a function of couplings  $\tau$ ). Then, due to our choice (10)-(12),  $P(E_1 \cdots E_M)$  factorizes.

Using Fourier representation for  $\delta$  function

$$P(E_1 \cdots E_M) = \int_{-i\infty}^{i\infty} \prod_{\alpha=1}^{2^N} \frac{dE_{\alpha}}{2\pi} \exp \left[ \sum_{\alpha=1}^M E_{\alpha} \hat{E}_{\alpha} - \sum_{\alpha} \hat{E}_{\alpha} \sum_{i_1 \cdots i_p} \tau_{i_1 \cdots i_p} \sigma_{i_1}^{\alpha} \cdots \sigma_{i_p}^{\alpha} \right] \quad (16)$$

Let us use standard representation

$$\exp(a\sigma) = \cosh(a) [1 + \tanh(a)\sigma], \quad (17)$$

where  $\sigma = \pm 1$ .

We can derive, expanding

$$\prod_{i_1 \cdots i_p} [\cosh(\hat{E}_1) + m \sinh(\hat{E}_1)] \prod_{\alpha=2}^M \cosh(\hat{E}_{\alpha}) [1 + \tanh(\hat{E}_{\alpha}) \sigma_{i_1}^{\alpha} \cdots \sigma_{i_p}^{\alpha}] \quad (18)$$

the expression for the  $P(E_1 \cdots E_M)$

$$P(E_1 \cdots E_M) = \int_{-i\infty}^{i\infty} \left\{ [\cosh(\hat{E}_1) + m \sinh(\hat{E}_1)] \cosh(E_1) \cdots \cosh(E_M) \right\}^{\alpha N} \prod_{\alpha=1}^M \frac{dE_{\alpha}}{2\pi} e^{\sum_{\alpha} E_{\alpha} \hat{E}_{\alpha}} \quad (19)$$

in the expression we neglect by effects of order

$$\sum_{1 \leq i_1 \cdots i_p \leq N} (\sigma_{i_1}^{\alpha} \sigma_{i_p}^B) \cdots (\sigma_{i_1}^{\alpha} \sigma_{i_p}^B) / C_N^P \quad (20)$$

We can consider this expression as a scalar product between 2 configurations).

Let us calculate (20) for the case, when only 1 turned spin. it is ease to derive

$$(C_{N-1}^P - C_{N-1}^{P-1}) / C_N^P = \frac{N - 2P}{N} \quad (21)$$

We see, that corrections disappear for the choice

$$P = \frac{N}{2} \quad (22)$$

When the number of turned spins is  $\delta k$ , we have expression like

$$\frac{1}{N} e^{-C\delta N} \quad (23)$$

In future calculations we using expression (19) for  $P(E_1 \dots E_M)$ . Then accuracy of one expression is polynomial or exponential (by  $N$ ). We call this corrections as "finite  $P$  corrections". Their investigation is a very hard work.

In the next section we calculating finite size effects for the model (15),(19) using the formula from [1]

$$\langle \ln Z \rangle = \Gamma'(1) - \int_{-\infty}^{\infty} \ln t e^{-\phi} d\phi \quad (24)$$

where

$$e^{-\phi} = \langle e^{-tZ} \rangle \quad (25)$$

It is easy to derive, taking  $Z = e^A$ , equation

$$A = \Gamma'(1) - \int_{-\infty}^{\infty} u d \left[ e^{-e^{U+A}} \right] \quad (26)$$

## 2 Paramagnetic Phase

In this phase, as well as in the spin-glass case, we taking  $m = 0$ .

It is easy to derive

$$e^{-\phi} = \left[ \int_{-\infty}^{\infty} \frac{dE_1 dE_2}{2\pi} e^{-E_1 E_2 + \alpha N \ln \cosh E_1 - t e^{-B E_2}} \right]^{2^N} \quad (27)$$

Let us make transformation

$$x = e^{-B E_2}, \quad E_1 = B E \quad (28)$$

After this easy to derive

$$\begin{aligned}
e^{-\phi} &= \left[ \int_{-i\infty}^{-i\infty} \frac{dE}{2\pi} \int_0^\infty e^{-x+(E-1)\ln x - UE + \alpha N \ln \cosh(BE)} \right]^{2^N} \\
&= \left[ \int_{-i\infty}^{-i\infty} \frac{dE}{2\pi} \Gamma(E) e^{-UE + \alpha N \ln \cosh(BE)} \right]^{2^N}
\end{aligned} \tag{29}$$

In this integral integration loop passed pole zero from right side.

For the integration we must lift integration loop righter, until saddle point, defined by equation

$$\tanh(BE) = \frac{U}{\alpha NB} \tag{30}$$

Doing this we must take care for contribution of intersected poles.

To calculate thermodynamic limit it is enough to take account poles  $0, -1$

$$e^{-\phi} \simeq \left[ 1 - e^{U + \alpha N \ln \cosh B} \right]^{2^N} \simeq \exp \left[ -e^{U + N \ln 2 + \alpha N \ln \cosh B} \right] \tag{31}$$

Using formula (24), we derive

$$\langle \ln Z \rangle \approx N \ln 2 + \alpha N \ln \cosh B \tag{32}$$

Let us take in to account finite size effect. The saddle point gives

$$\begin{aligned}
e^{-\phi} &\simeq \left[ 1 - e^{U + \alpha N \ln \cosh B} + \Gamma(E) e^{-EU + \alpha N \ln \cosh(BE)} \right]^{2^N} \\
&\simeq \left[ e - e^{U + \alpha N \ln \cosh B} + \Gamma(E) e^{-(E+1)U + \alpha N \ln \cosh(BE) \cosh B} \right]^{2^N} \\
&= e^{-U + \alpha N \ln \cosh B + N \ln 2 + \varphi(U)}
\end{aligned} \tag{33}$$

So we found connection between  $\varphi$  and  $U \equiv \ln t$

$$\ln \varphi = U + U_0 + \varphi(U) \tag{34}$$

where

$$\begin{aligned}
U_0 &= -(N \ln 2 + \alpha N \ln \cosh B) \\
\varphi(U) &= \Gamma(E) \exp \left[ -(E+1)U + \ln \cosh(BE) / \cosh B \right]
\end{aligned} \tag{35}$$

Let us express  $U$  by  $\ln \phi$

$$u \approx -\ln \phi - U_0 - \varphi(U_0) \quad (36)$$

With such accuracy

$$\begin{aligned} \langle \ln Z \rangle &= \int_0^\infty \ln t (e^\phi d\phi) \\ &= \alpha N \ln \cosh B + N \ln 2 + \Gamma(E) e^{N(E+1) \ln \delta + \alpha \ln [\cosh(BE) (\cosh B)^E]} \end{aligned} \quad (37)$$

In the last expression we have neglected preexponent besides  $\Gamma(E)$ . The next subphase appears, when saddle point  $E$  coincides with point-2:

$$\tanh(2B) = \frac{\alpha \ln \cosh B + \ln 2}{\alpha B} \quad (38)$$

The equation about subphase of paramagnetic phase demands more careful discussion.

### 3 Spin-glass phase

In this case it is enough to intersect only one pole  $E = 0$

After transformation we have

$$e^{-\phi} = \left[ 1 + \frac{1}{2\pi} \int_{-i\infty}^{i\infty+E_0} dE \Gamma(E) e^{-EU + \alpha N \ln \cosh(BE)} \right]^{2^N} \quad (39)$$

$$e^{-\phi} \equiv \exp \left[ \frac{-|\Gamma(E_0)|}{h\sqrt{2\pi}} \exp(-E_0 U + \alpha N \ln \cosh(BE_0) + N \ln 2) \right] \quad (40)$$

where  $h = \sqrt{\alpha N B}$  and for  $E_0$  we have equation

$$\tanh(BE_0) = U/\alpha N B \quad (41)$$

Let us express again  $U$  by  $\ln \phi$

$$\ln \phi \approx E_0 U + \alpha N \ln \cosh(BE_0) + N \ln 2 - \frac{1}{2} \ln N \quad (42)$$

$$U \approx \frac{1}{E_0} \ln \phi + \frac{\alpha N}{E_0} \ln \cosh(BE_0) + \frac{N \ln 2}{E_0} - \frac{1}{2} \frac{\ln N}{E_0} \quad (43)$$

In this formula we deal with expressions like  $BE_0$  and  $U/B$ .

$$\tanh(BE_0) = \frac{\ln \cosh BE_0 + \ln 2/\alpha}{BE_0} - \frac{\ln N}{2\alpha N BE_0} + \frac{\ln \phi}{\alpha N BE_0} \quad (44)$$

If we neglecting the last terms in (44) then we have

$$E_0 = \frac{-B_c}{B} \quad (45)$$

Where at  $B_c$  disappears entropy. For corrections we have ( $x = BE_0$ )

$$\begin{aligned} (x \tanh x - \ln \cosh x - \ln 2\alpha) &= -\frac{\ln N}{2\alpha N} \\ x &= x_0 - \frac{(\cosh x_0)^2 \ln N}{x_0 2\alpha N} \\ U = \alpha N B \tanh x &= \alpha N B \tanh x_0 - \frac{1}{2} \ln N \frac{B}{B_c} \end{aligned} \quad (46)$$

For the free energy we receive

$$\langle \ln Z \rangle = \alpha N B \tanh B_c - \frac{1}{2} \frac{B}{B_c} \ln N \quad (47)$$

So at phase transition point free energy has jump

$$-\frac{1}{2} \ln N \quad (48)$$

as for the case fully connected model

## 4 Ferromagnetic Phase

For  $e^{-\phi}$  we have expression

$$e^{-\phi} \approx G(U) \tilde{G}(U)^{2^N - 1}$$

where

$$\tilde{G}(U) = \begin{cases} 1 - \frac{\cosh BE_0 |\Gamma(E_0)|}{2\sqrt{\pi}} e^{-E_0 U + \alpha N \ln \cosh(BE_0)} & , E_0 < 0; \\ \frac{\cosh BE_0 |\Gamma(E_0)|}{2\sqrt{\pi}} e^{-E_0 U + \alpha N \ln \cosh(BE_0)} & , E_0 > 0 \end{cases} \quad (49)$$

and for  $E_0$  we have equation

$$\tanh(BE_0) = \frac{U}{\alpha N B} \quad (50)$$



Similarly for  $G(U)$

$$G(U) = \begin{cases} 1 - \frac{(\cosh(BE_1) - m \sinh(BE_1))}{2\sqrt{\pi}\sqrt{1-m^2}} |\Gamma(E_1)| e^{-E_1 U + \alpha N \ln[\cosh(BE_1) - m \sinh(BE_1)]} & , E_1 < 0 \\ \frac{\cosh BE_1 - m \sinh BE_1}{2\sqrt{\pi}\sqrt{1-m^2}} |\Gamma(E_1)| e^{-E_1 U + \alpha N \ln[\cosh(BE_1) - m \sinh(BE_1)]} & , E_1 > 0. \end{cases} \quad (51)$$

and for  $E_1$

$$\frac{m \tanh BE_1 - m}{1 - m \tanh E_1} = x, \quad x = \frac{U}{\alpha NB} \quad (52)$$

Let us define function

$$f(x) = \frac{1+x}{2} \ln \frac{1+x}{1+m} + \frac{1-x}{2} \ln \frac{1-x}{1+m} \quad (53)$$

$$g(x) = \frac{1+x}{2} \ln \frac{1+x}{2} + \frac{1-x}{2} \ln \frac{1-x}{2} + \ln 2 \quad (54)$$

$$(55)$$

Simple calculations give

$$-E_0 U + \alpha N \ln \cosh(BE_0) = -\alpha N g(x) \quad (56)$$

$$-E_1 U + \alpha N \ln \cosh(BE_1) = -\alpha N f(x) \quad (57)$$

We want to calculate corrections

$$\langle \ln Z \rangle - \alpha NBm = \int G(U) [1 - \tilde{G}(U)^{2^N-1}] \equiv I \quad (58)$$

It is convenient to break integration interval

$$I = \int_{-\infty}^{-U_0} + \int_{-U_1}^{-U_1} + \int_{-U_1}^{\infty} \quad (59)$$

Where

$$U_0 = \alpha NBm \quad (60)$$

and

$$U_1 = \alpha NBx_1 \quad (61)$$

The value of  $x_1$  defined from equation

$$\alpha g x_1 = \ln 2 \quad (62)$$

In the first integral we see the condition of dominance of ferromagnetic free energy under spin-glass expression:

$$\alpha g(m) > \ln 2 \quad (63)$$

The main contribution to the finite size correction comes from second integral.

We have two subphases. In the first out to saddle point is in the integral of integration, in the second one-out side.

$$I \sim \exp[-\alpha N f(-x_1)], \quad x_1 > x_2 \quad (64)$$

and

$$I \sim \exp[-\alpha N f(-x_2) - \alpha N g(x_2) + N \ln 2], \quad x_1 > x_2 \quad (65)$$

where for  $x_2$  we have

$$x_2 = \frac{1 - \sqrt{1 - m^2}}{m} \quad (66)$$

Our expression (64),(65) coincide with known results of information theory

## 5 Summary

We have calculated finite size effects for Derrida model with weak connectivity.

In the spin-glass phase we received logarithmic corrections, in paramagnetic-exponential.

In ferromagnetic phase at  $T = 0$  our results coincide with information theory (the results of random coding). That codes give for the value of connectivity near the critical extremal possible suppression of decoding error probability. For strong connectivity the result is unknown for optimal coding.

Unfortunately known boundaries from information theory show, that Derrida's model does not give extremal suppression for the case of high values of connectivity (weak velocity). To clarify situation, we need consider finite  $P$  corrections.

In the work [10],[11] has suggested to consider

$$\langle |\sigma_i(\tau, T_N)| \rangle_\tau \quad (67)$$

instead of magnetization at  $T = 0$ , where  $T_N = \frac{1}{2} \ln \frac{1+m}{1-m}$

Neglecting finite  $P$  corrections we received the same result for the exponent, as for the case  $T = 0$

We would like to thank Yev.Harutunyan for consultations during the work.

We are also grateful L. Basalyga, R.Dobrushin, B.Derrida, Sh.Rouhani. N.Sourlas for useful remarks.

This work was supported by NATO linkage grant LG 9303057, ISF grant MVMOOO and German ministry of research and technology grant 211-5231. D.B. Saakian thanks Ivan Kostov for hospitality in Saclay.

## References

- [1] B.Derrida,Phys.Rev.Lett.45(1980),79.
- [2] D.Gross, M.Mezard,Nuci.Phys.,**B240**(1984) 43.
- [3] N.Sourlas, Nature 339,(1989),693
- [4] D. B.Saakian, JETP Lett.(1992),**56**, n.5
- [5] D. B.Saakian, Phys.Lett.(1993),**A180**,169
- [6] I.Chisar, J.Korner, Information theory,Moscow,(1985)
- [7] D. B.Saakian, JETP Lett.(1995),**56** n.5.
- [8] D. B.Saakian, Sh.Rouhani, .Mod.Phys.Lett. B.(1995)
- [9] C.Dominicis, P.Mottishaw, J.Phys.(1987),**A20**, 1267.
- [10] P.Rujan, Phys.Rev.Lett.(1993),**70** 2961

[11] H.Nishimori,Phys.,(1994),**A204**, 1.

## Appendix A

In this appendix we show, how transform our results for the case of full connectivity and derive results of [1].

For this purposes we introduce coupling  $J > 0$ , so we consider

$$\begin{aligned}
P(j_{i_1 \dots i_p}) &= \left(1 - \frac{\alpha N}{2C_N^p}\right) \delta(j_{i_1 \dots i_p}) \\
&+ \frac{\alpha N}{2C_N^p} \left[ \frac{1+m}{2} \delta(j_{i_1 \dots i_p} - J) + \frac{1-m}{2} \delta(j_{i_1 \dots i_p} + J) \right]
\end{aligned} \tag{A.1}$$

For this purposes it is just enough to replace  $B \rightarrow BJ$ ,  $B_c \rightarrow B_c J$

It is easy to check [6], that transition to full connectivity means

$$J \rightarrow J\alpha^{-1/2}, \quad \alpha \rightarrow \infty \tag{A.2}$$

For us it means

$$e^{-\phi} = \left[ \int_{-i\infty}^{i\infty} \frac{dE}{2\pi} \Gamma(E) e^{-Eu + \frac{N(JBE)^2}{4}} \right]^{2N} \tag{A.3}$$

We consider the case  $m = 0$ .

Different expressions for  $e^{-\phi}$  depends on the poles, intersected by our loop for saddle point intergation at  $E = 2U/\lambda^2$  and the last depends on the temperature. It is easy to derive (A.3) directly.